

# Mild head injury and concussion



the brain injury association

This publication is part of Headway's *About the brain* series. To browse through our publications on a range of issues relating to brain injury and download these free-of-charge, visit [www.headway.org.uk/information-library](http://www.headway.org.uk/information-library).

## Contents

Introduction .....	2
Section one: What is a mild head injury? .....	2
Initial assessment .....	3
Early stages after discharge .....	3
Dos and don'ts in the first few days after injury .....	5
Section two: Post-concussion syndrome .....	6
Common symptoms and practical suggestions .....	6
Cognitive symptoms .....	7
Physical symptoms .....	10
Emotional and behavioural symptoms .....	18
Section three: Other issues .....	21

This publication is also available as a printed booklet.  
For more information or to order, contact 0115 924 0800  
or visit [www.shop.headway.org.uk](http://www.shop.headway.org.uk).

People directly affected by brain injury can receive limited free copies of Headway print booklets by contacting the helpline on 0808 800 2244.

\*print copy may contain minor differences due to revision of content



This publication has been kindly sponsored by:



**Cycle-SOS**  
SIMPLY OUTSTANDING SOLICITORS

ADVERTISEMENT

**Providing specialist legal services to injured cyclists**

Freephone 0808 100 9995    [cycle-sos.co.uk](http://cycle-sos.co.uk)

Proud supporters of Headway – the brain injury association

---

## Introduction

Head injuries come in a variety of guises. Some are severe, with loss of consciousness lasting days or weeks and can cause significant neurological disabilities afterwards. Fortunately, most are classified as mild. However, even so-called 'mild' head injuries are associated with physical, cognitive and emotional symptoms.

This publication is written for anyone who has had what is classified as a mild head injury and for their family and friends. While the majority of people recover from a milder head injury very quickly, experiencing one can still cause feelings of worry and concern. However, understanding the symptoms, knowing what to look out for and learning how to cope with the effects can help at a time that, for some, can be scary and confusing.

---

## Section one: What is a mild head injury?

Mild head injury is also often referred to as concussion, minor head injury or mild traumatic brain injury (mTBI). Regardless of the terminology used, the occurrence of a head injury in these cases causes the brain to shake back and forth inside the skull, causing mild damage. While most mild head injuries result in no long-term damage to the brain, it can cause temporary disruption to brain function that can last for at least a number of weeks.

It is estimated that over a million people each year attend Accident and Emergency (A&E) departments in the UK after a head injury, with the majority of these injuries being classed as mild. Mild head injury is commonly caused by falls, road traffic collisions, assaults and sports accidents (information about mild head injuries sustained specifically through sporting accidents is available in our publication [\*Concussion in sport\*](#)).

Mild head injuries can also be sustained during relatively common accidents such as hitting one's head against a wall or an object dropping onto the head.

There are currently a number of ways of classifying mild head injury, but at

present it is most widely defined by:

- Loss of consciousness of less than 30 minutes (or no loss of consciousness)
- Post-traumatic amnesia (PTA) of less than 24 hours after injury. This is a period where people are confused, act strangely and are unable to remember what has just happened. More information on PTA is available in our publication [Post-traumatic amnesia](#).

It is important to note that only around 10% of reported mild head injuries involve a loss of consciousness - so it is important to not solely rely on this as an indicator.

## Initial assessment

Patients who are suspected of having a head injury should be assessed using the Glasgow Coma Scale (GCS), which assesses their level of consciousness on the basis of neurological signs and responsiveness. The scale ranges from 3 - 15 with 3 being seen in death or deep coma, and 15 associated with being fully conscious and not confused. For a mild head injury, the initial score will have been 13-15 and patients shouldn't be discharged until their GCS is 15.

If patients are considered to be in danger of complications, they will be given a CT scan when they are initially seen in hospital. CT stands for computerised tomography. A series of X-rays are taken to build up a detailed picture of the inside of the skull and brain. CT scans of the head are an effective method of checking for a possible brain haemorrhage (bleeding in or around the brain) or swelling of the brain. In some cases it may be necessary to perform an MRI scan or other investigations after the CT scan in order to get more detailed information.

For more information about scans commonly used to detect brain injury, see our publication [Scans and tests after brain injury](#).

## Early stages after discharge

In the early days of a mild head injury, it is common to feel symptoms such as dizziness, tiredness, headaches and problems with your memory, among other

Helpline: 0808 800 2244  
Email: [helpline@headway.org.uk](mailto:helpline@headway.org.uk)  
Web: [www.headway.org.uk](http://www.headway.org.uk)

symptoms. The section of this publication 'Common symptoms and practical suggestions' discusses these in more detail.

Although it is unlikely, serious complications can sometimes arise in the days and even weeks after a mild head injury. It is important that you, and a responsible adult who can monitor your condition, are aware of the symptoms listed below which could indicate a more serious injury:

- Any loss of consciousness
- Drowsiness or deteriorating consciousness
- Loss of memory for events before or after the injury
- Confusion or irritability out of character, or unusual behaviour
- Problems with understanding information or communicating
- Loss of balance or problems with walking
- Decreased sensation, or pins and needles down arms/ legs
- Blurred or double vision
- Weakness
- Any fits (collapsing or passing out suddenly)
- Severe and increasing headache not relieved by painkillers such as paracetamol
- Repeated vomiting
- Reduced neck movement or severe neck pain
- Deafness in one or both ears that was not present before
- Clear fluid coming out of the ears or nose
- Inability to be woken
- Bleeding from one or both ears

**If you experience any of these symptoms, it is important that you go to a hospital Accident and Emergency department immediately.**

Helpline: 0808 800 2244  
Email: [helpline@headway.org.uk](mailto:helpline@headway.org.uk)  
Web: [www.headway.org.uk](http://www.headway.org.uk)

In the first 24-48 hours of the concussion, it is important to rest and sleep as needed to give the brain time to recover. 'Resting' includes reducing screen time, such as watching TV or scrolling through social media. The person should also reduce stress where possible and avoid situations where they may feel overstimulated, for example being in busy environments. Easy activities of daily living and gentle exercise such as walking can continue, and can be helpful with recovery. Once symptoms have reduced, the person can progress through the graduated return to activity (education/ work) and sport guidance produced by the UK Government and Sport + Recreation Alliance (SRA), available at [www.sportandrecreation.org.uk/policy/research-publications/concussion-guidelines](http://www.sportandrecreation.org.uk/policy/research-publications/concussion-guidelines).

Further suggestions of what to do in the weeks that follow are...

### **Do...**

- Do get rest and gentle exercise such as walking for the first 24-48 hours of the concussion being sustained.
- Avoid intense exercise.
- Do stay within reach of a telephone in case you need medical help over the next few days
- Do share this information with a relative or friend who can keep an eye on your condition.
- Do follow the UK government/ SRA returning to activity (education/ work) and sport guidance

### **Don't...**

- Don't stay at home alone for 24 hours after the injury
- Don't play or train for contact sports for at least 14 days without consulting an appropriate healthcare professional
- Don't drink alcohol for 24 hours after injury and until symptoms improve
- Don't return to driving for 24 hours after injury and until your symptoms

Helpline: 0808 800 2244  
Email: [helpline@headway.org.uk](mailto:helpline@headway.org.uk)  
Web: [www.headway.org.uk](http://www.headway.org.uk)

improve. If in doubt, consult your GP. Commercial drivers should be reviewed by an appropriate medical professional before returning to driving.

---

## **Section two: Post-concussion syndrome**

A wide variety of physical, cognitive and emotional symptoms can be seen after a mild head injury. The symptoms after a head injury can be very distressing, especially if you don't know what to expect. In most cases, these symptoms resolve themselves within a few days or weeks, though they may persist for a much longer period of time. Post-concussion syndrome, also known as post-concussive syndrome, is the name given to the constellation of symptoms which continue to occur following a mild head injury.

These continuing symptoms are likely to resolve themselves within a few days or weeks. However, in a minority of cases, problems may persist for longer, but still resolve themselves eventually. This can be a frustrating and distressing time, as the symptoms may be subtle, and you may not feel that you are back to a sense of full health.

This may also be the most prolonged period of feeling ill that you have experienced and you may wonder if you will ever feel better. Following the suggestions in this section should help to make you feel better as quickly as possible. The most important thing to remember is to be patient and kind to yourself and try not to rush things.

If you have these symptoms for more than a few weeks after the injury, or if they are severe, worsening and not getting any better, then you should see your GP. Your GP might be able to refer you to a neurologist, neuropsychologist or neuropsychiatrist for further support.

### **Common symptoms and practical suggestions**

The symptoms outlined in this section are separated into cognitive symptoms, physical symptoms, and emotional and behavioural symptoms.

Helpline: 0808 800 2244  
Email: [helpline@headway.org.uk](mailto:helpline@headway.org.uk)  
Web: [www.headway.org.uk](http://www.headway.org.uk)

You might experience any number of these symptoms at different times and the effects of a mild head injury are different for everyone. This section offers information on the most common symptoms and practical suggestions to help you recover more quickly. It may be possible to treat each one of the symptoms in isolation, so it is important to consult your GP about possible referral to a relevant specialist.

This section contains references to other Headway publications which may be of help. Headway publications are available to download free-of-charge from [www.headway.org.uk/information-library](http://www.headway.org.uk/information-library) or by contacting the Headway helpline on 0808 800 2244 or [helpline@headway.org.uk](mailto:helpline@headway.org.uk).

## Cognitive symptoms

Cognitive symptoms are very commonly experienced after a mild head injury. These symptoms can take the form of problems with memory, concentration and organisation. They can also be quite subtle. Stress and fatigue can worsen cognitive symptoms, so it is very important to practice self-care.

### *Memory problems*

Loss of memory can make everyday life frustrating. You may be able to remember your wedding day, but not where you put your shoes. This can be because you're having problems concentrating and also because mild head injury disrupts the memory system for a while. Your memory should improve as you recover, but in the meantime there are ways that you can help yourself.

- Use calendars and diaries to remind you of events
- Use stick-on labels as reminders, for example, to make sure you lock the back door
- Carry a notebook to write down new information and things you need to remember
- Use cooking timers in the kitchen

Helpline: 0808 800 2244  
Email: [helpline@headway.org.uk](mailto:helpline@headway.org.uk)  
Web: [www.headway.org.uk](http://www.headway.org.uk)



- Assign a specific place for everyday items and make sure you keep them there
- Use calendars and reminder apps on mobile phones. However, try to avoid overusing such electronic devices, as these can be quite cognitively demanding and can affect your quality of sleep.
- Break down information into small chunks to make it easier to remember
- Get plenty of sleep and rest; your memory will work better if you are fresh and rested

For more information on this subject, see our publication [Memory problems after brain injury](#).

### *Difficulties with attention and concentration*

The ability to concentrate and attend to things is very often significantly reduced by mild head injury, particularly in the first few days. The problem usually takes two forms.

Firstly, it may be difficult to concentrate on more than one task, or to concentrate when there are distractions, such as when children are playing or the TV is on. You may find it difficult to cope in large groups of people when several conversations are going on at once.

Secondly, the concentration span can be reduced so you may find it difficult to concentrate on one thing for as long as usual. This can be particularly disruptive to work and study.

Some suggestions for dealing with attention and concentration problems are as follows:

- Start a task when you are feeling fresh, then stop as your attention begins to fade
- Cut out background noise
- Try to carry out important or complex tasks in a quiet area
- Do one task at a time

Helpline: 0808 800 2244  
Email: [helpline@headway.org.uk](mailto:helpline@headway.org.uk)  
Web: [www.headway.org.uk](http://www.headway.org.uk)



- Break down tasks into manageable chunks
- Concentration depends on being alert, so if you are tired and can't concentrate, have a rest

### *Difficulties with planning and organising*

You may find it difficult to plan and organise. This can be frustrating, especially if these skills are needed for your job, and even everyday tasks such as preparing meals can become difficult. Some suggestions to help compensate for these difficulties are:

- Try to stick to a daily routine
- If your job requires high levels of planning and organisation, consider returning on part-time hours and taking on lighter duties until you feel better. Remember not to rush into returning to full-time work too soon, as this can result in a loss of confidence and can lengthen the period of recovery.
- Try to stick to simple meal plans - follow set recipes from cookery books and, if possible, get someone to help
- Make 'to do' lists
- Set realistic goals for each day and don't be too disheartened if you don't always achieve them
- Use external aids, such as mobile phones, pagers, electronic watches or tablets, to provide reminders of tasks
- Use a diary, journal or online calendar (e.g. Google Calendar)
- Do one task at a time
- Try to plan and carry out tasks in a quiet area
- Get plenty of rest
- Accept that recovery is often a journey and don't feel bad about yourself if you make mistakes

For more information on this subject, see our publications [\*Decision making difficulties after brain injury\*](#) and [\*Executive dysfunction after brain injury\*](#).

## Physical symptoms

### *Sleep problems*

People tend to need more sleep than usual for a while after a mild head injury. Getting enough sleep and maintaining good sleep habits are essential to a good recovery. Lack of sleep leads to fatigue, which can worsen a lot of other problems, such as difficulty concentrating, headaches and irritability. The following dos and don'ts should help you to maintain good sleep habits:

- Do establish a regular routine by going to bed and getting up at the same times each day
- Do take 'power naps' during the day but try to avoid doing so after 4pm, as this may disrupt your sleep/wake cycle
- Do use an alarm clock to wake you from naps to ensure you don't sleep too long
- Do use your bed for sleeping only – don't watch TV in bed
- Do develop rituals before bedtime to wind down and relax; for example, taking a bath or listening to gentle music
- Do get regular exposure to the outdoors and natural light
- Do exercise regularly
- Do create a calm bedroom that is cool, dark and quiet – earplugs and blackout blinds can help; remember that the light from electronic devices such as phones and tablets can affect sleep cycles, so switch these off if possible
- Do avoid stress and worry at bedtime
- Do use relaxation techniques such as mindfulness

- Don't push yourself to stay awake and don't worry about 'giving in' to tiredness
- Don't eat heavy meals late in the evening
- Don't look at the time when you wake up as this may make it difficult to get back to sleep
- Don't have too much caffeine, nicotine or alcohol in the evening

As you recover you will find that you need less sleep. However, you may find that you can't get to sleep, or wake during the night. Typically, this can be made worse by:

- Stress – This may be caused by frustration with your rate of recovery, or by other people expecting you to recover sooner. Financial or family problems may also contribute, particularly if you are still absent from work.
- Changes in your sleep cycle – If you don't have regular times for getting up and going to bed then your sleep might be affected.
- Depression – This can affect your sleep cycle by making it difficult for you to get to sleep at night and making you wake up early in the morning.

At this stage there are a number of strategies that can help:

- Relaxation training. Ask your GP about techniques such as meditation and yoga. You could also ask about referral to a counsellor who can teach relaxation techniques.
- If you have specific worries that are affecting your sleep try to resolve them. Talk the matter through with a close friend or family member, or ask your GP about referral to a counsellor.
- If you have cut out your daytime nap and are feeling more tired, you could try taking naps again and then gradually phasing them out. If you are still taking daytime naps and are having trouble getting to sleep at night, try cutting out the naps.

## *Fatigue*

This is one of the most common effects of mild head injury and it is also the most important symptom to control, as being tired affects everything you do. You may get tired more quickly than before and even simple actions, like dressing or cooking, can make you feel exhausted. Following the suggestions in the previous section on sleep problems should help you to manage your fatigue and there are a number of other things you could try:

- Set a daily routine – prioritise and set realistic targets to achieve each day
- Pace your workload by taking lots of short breaks during tasks
- Think about when you are most tired – try to do your more difficult and demanding tasks at times when you are most awake
- Identify the triggers to your fatigue – if you can recognise the activities and situations that are most tiring you can plan for them in your daily routine
- Build rest times into your day
- Don't overdo it, but try to build activities up slowly over time
- Explain to other people that you get tired – it can be difficult to recognise the signs of fatigue after a head injury, so ask family and friends to let you know when you appear fatigued and the signs they notice
- If going back to work, start with fewer hours and build up gradually

Don't push yourself or you may get into a cycle where you don't get enough rest. In the space of a few days you could become completely exhausted and all other symptoms may get worse. This is referred to as 'boom and bust' and you should take care to avoid it. Your fatigue will decrease as you start to feel better but can return, along with other symptoms, if you try to do too much too soon. For further information on fatigue and sleep you may find it useful to refer to our publication [\*Fatigue after brain injury\*](#).

## *Headaches*

Headaches are a common complaint after mild head injury, particularly in the early stages. Remember, if the headache is severe, persistent and unrelieved by

ordinary painkillers in the few days after injury then it can be a sign of complications and you should return to your nearest A&E department.

In the later stages, headaches are often caused by other difficulties. For example, if you are fatigued, stressed, physically tense, or struggling with your concentration, you may get headaches. At this stage the following suggestions may help:

- Write down when you get headaches – you may be able to see if there is a time or place that causes them and plan for that
- Try relaxation techniques – often headaches are due to muscle tension and techniques such as meditation and even moderate exercise could help
- Try not to take too many painkillers (especially ones containing codeine) as these can become addictive and can actually make headaches worse in the long run
- You should discuss these issues with your GP or other healthcare professional

For more information on this topic, see our publication [Headaches after brain injury](#).

### *Dizziness, balance and co-ordination problems*

Mild head injury often disrupts the balance organs in the ears, so it is common in the first few days after injury to experience vertigo, where the world seems to spin around you. The slowing of information processing caused by the injury can also lead to a feeling of unreality, disorientation and a sense of ‘floating’.

Symptoms which are closely related to dizziness are problems with balance and co-ordination. You may feel wobbly and unsteady on your feet because of the dizziness. You may also drop things or bump into people and objects due to your brain processing information less efficiently than usual.

You can be assured that these problems are common and usually settle within a few days or weeks and then disappear entirely. Take greater care in situations which could be dangerous, such as crossing the road, and don’t drive until the

symptoms have gone. If the symptoms are very problematic or persistent then see your GP as there are some therapeutic techniques which may help. For more information on this, see our publication [\*Balance issues and dizziness after brain injury\*](#).

### *Visual problems*

It is important to remember that if you experience blurred or double vision in the first few days after the injury you should return to your nearest A&E department. If you still experience double vision in the longer term then legally you must not drive and should consult your GP about this (for further guidance on this issue, see our publication [\*Driving after brain injury\*](#)). Some other problems with vision that are common after mild head injury are:

- Loss of depth perception
- Photophobia (sensitivity to light)

These symptoms are usually mild and should clear up within a week or two. However, if they persist or are very debilitating then see your GP. Be extra careful in situations where clear, accurate vision is required and avoid driving until the symptoms have cleared. If your depth perception is affected, be very careful when crossing the road. Sensitivity to light is one of the most common visual problems after mild head injury. This can even be triggered by indoor lighting, especially fluorescent lights and it can help to wear sunglasses or tinted glasses, even indoors. This problem can remain when other symptoms have cleared, meaning sometimes people have to carry on wearing dark glasses, even when they have otherwise recovered.

For more information on this issue, see our publication [\*Visual problems after brain injury\*](#).

### *Hearing problems*

Hearing problems usually take the following forms:

- Mild deafness
- Sensitivity to loud noise

Helpline: 0808 800 2244  
Email: [helpline@headway.org.uk](mailto:helpline@headway.org.uk)  
Web: [www.headway.org.uk](http://www.headway.org.uk)

- Tinnitus - the perception of sound without external cause. This can take the form of ringing, buzzing, humming, whistling, whining or other noises.

Any deafness could be caused by damage to the inner ear or nerves in the brain and should be reported to a doctor to rule these possibilities out. Alternatively, you may only notice the impaired hearing when you are in an environment such as a party, when many conversations are going on at once. In this case the impaired hearing may be caused by reduced attention and concentration and should get better as information processing ability gets back to normal.

Sensitivity to loud noise is common after mild head injury. As with other symptoms this will go away in time, but until it does the only real strategy is to avoid noisy atmospheres that you find distressing. Tell family and friends about the problem and ask them to respect your need for a quiet environment.

Avoiding noisy situations can lead to avoiding environments that you would usually enjoy, such as parties and football matches. Talk to family and friends about this and try to organise enjoyable, social occasions where you can control the noise levels. For example, ask a select few friends around for a quiet dinner.

Tinnitus is not one of the most common effects of mild head injury, but can be extremely distressing when it occurs. See a doctor for an examination to rule out possible treatable causes and ask about referral to an ear specialist.

### *Communication problems*

You may find it difficult to express yourself in an accurate way. You may not be able to find the right words, or you might muddle sentences. The following suggestions may help:

- Be patient with yourself
- Take the time to find the right words
- Prepare what you want to say by thinking of key words or making lists
- Explain to people that they need to be patient with you



You can also ask your GP to refer you to a Speech and Language Therapist who could be able to help. For further information on this subject see our publication [Communication problems after brain injury](#).

### *Changes to taste or smell*

Following a mild head injury some people report that their senses of taste and/or smell have been affected. The senses of taste and smell are linked so if the sense of smell is lost then the sense of taste will also be affected.

Changes to the senses of taste and smell can be extremely distressing and sadly there are no cures available. Recovery often occurs spontaneously after mild head injury, often within the first three months. In the meantime, there are strategies you can use to compensate and stay safe:

- Fit a smoke alarm, have electrical appliances regularly serviced, unplug appliances when not in use and use an alarm to remind you of food cooking in the oven
- Have gas appliances regularly serviced and fit a gas detector; you might want to consider fitting an electric cooker and fire
- Always eat or throw out food by its 'use by' date; clear out the fridge and cupboards regularly
- Try to keep products such as drinks, bleach, cleaning chemicals and solvents in their original bottles/packets; make sure they are clearly labelled
- Empty rubbish bins regularly and keep toilets and kitchen appliances clean to avoid health risks
- Be aware of the need to wash yourself, your clothes and bed-sheets regularly; use an antiperspirant deodorant and perhaps a shoe deodoriser too
- Ask a close family member, loved one or friend to let you know if your body odour is an issue
- It is important to keep your mouth clean, brush teeth regularly and

thoroughly, and pay regular visits to the dentist; using mouthwash and dental floss helps

- Take precautions and follow manufacturer's advice when using products such as paint, cleaning chemicals and solvents; wear a protective mask, ensure rooms are well ventilated and don't smoke

It is also important to remember to maintain a healthy, balanced diet, as loss of taste and smell can affect eating habits.

For further information on this subject see our publication [\*Loss of taste and smell after brain injury\*](#). You can also find information on healthy eating on the British Dietetic Association website at [www.bda.uk.com](http://www.bda.uk.com) and on the NHS Choices website at [www.nhs.uk/healthyeating](http://www.nhs.uk/healthyeating).

### *Sexual problems*

You may find that you have little or no sexual urge, or the complete opposite. In some cases this can continue longer than other symptoms and can be a major factor in preventing a return to a full life.

If you have a partner, it is important to speak to them about this, as it can make relationships difficult. If you don't have a partner, it is also important to discuss this with someone, as it can affect the way you feel about yourself and can lead to problems with depression.

The problems may be caused by several factors. They may be the direct result of injury to the brain, but after a mild head injury, they are more likely to be a result of the psychological effects of other symptoms.

Alternatively, sexual problems can be a result of damage to the pituitary gland, which is situated at the base of the brain and controls hormone secretion in the body. Further information specifically on hormonal imbalances is available in our publication [\*Hormonal imbalances after brain injury\*](#).

The following suggestions may help:

- Seek advice from your GP

Helpline: 0808 800 2244  
Email: [helpline@headway.org.uk](mailto:helpline@headway.org.uk)  
Web: [www.headway.org.uk](http://www.headway.org.uk)

- Seek referral to a clinical psychologist or neuropsychologist who specialises in sexual difficulties after head injury
- Speak to a counsellor or a specialist organisation such as [Relate](#), who deal with sex and relationship issues
- If there are possible hormonal problems, then discuss it with your GP and ask about referral to an endocrinologist

More information on sexual problems and tips for managing these are available in our publication [Sex and sexuality after brain injury](#).

### *Nausea*

It is common to experience a sensation of nausea after a mild head injury, especially in the first few days. If the nausea is severe or accompanied by vomiting, then it is important to seek medical advice. However, although it can be very unpleasant, usually the symptom isn't serious, is relatively mild and goes away in time. Nausea may be a result of the injury itself or could be caused by medication you're taking. As with other symptoms, if the problem persists it would be a good idea to discuss it with a GP.

## **Emotional and behavioural symptoms**

### *Irritability and anger*

Things that wouldn't bother you before your head injury may anger or irritate you. You might lose your temper for no reason or snap at family and friends. This happens because self-control and calmness require the brain to be fresh and working well. As you recover the problem should improve, but it can be very upsetting to you and those around you, so it is important to try to manage it.

- Fatigue and lack of sleep can cause irritability – if you are feeling very tired follow the suggestions in the 'Sleep problems' and 'Fatigue' sections of this publication
- A daily routine will make the causes of irritability easier to recognise – try to spot what activities contribute to irritability and anger and what times of day

Helpline: 0808 800 2244  
Email: [helpline@headway.org.uk](mailto:helpline@headway.org.uk)  
Web: [www.headway.org.uk](http://www.headway.org.uk)

the problem is at its worst

- Find alternative, healthy outlets for anger, for example, exercising at the gym
- Try relaxation techniques such as meditation, or just make sure you make time in your day to relax in whatever way you find most comfortable, such as listening to music, reading or having a long bath

If you find yourself in a situation where you recognise you may be about to lose control, take time out by leaving the room and taking some deep breaths.

For further information on this subject, see our publication [Anger after brain injury](#).

### *Depression and anxiety*

These problems are common after mild head injury and are often caused by other difficulties, such as:

- Tiredness
- Concentration problems
- Difficulty controlling emotions
- Symptoms persisting longer than expected
- Financial concerns
- Family and relationship problems

Worrying about your health can be very stressful and this may be the longest period you have ever felt ill. If the symptoms don't clear up as quickly as you had hoped, then depression and anxiety can become more of an issue.

One factor that can lead to depression is a lack of information on your condition, which makes some people think they are 'going mad'. Be assured that your symptoms are perfectly normal and should clear even if it takes longer than you expect.

Some important things to remember in order to help yourself are:

- Try to maintain good sleep habits – depression and anxiety lead to

interrupted sleep and the resulting tiredness makes the problems worse

- Try relaxation techniques such as meditation
- Write down your worries and work through them one at a time
- Try to think positively and don't dwell too much on negative things – maintaining a positive outlook can significantly improve your rate of recovery
- Talk through your worries with family, friends, a support group or GP
- If you have thoughts that life is not worth living or have thoughts of suicide, do seek help immediately and confide in a loved one, GP or if thoughts are particularly strong, attend A&E. It's not uncommon that people can feel despair after a significant life event such as a head injury. Seeking help and talking about it can be lifesaving.

If your problems persist and you are worried about your state of mind, see your GP. Make sure the GP understands that your depression stems from your injury and its effects, and ask about referral to a neuropsychologist, neuropsychiatrist or a counsellor who specialises in depression after head injury. It is particularly important to seek professional advice if you have a history of depression from before the injury, as this may make it more likely that you will experience depression now.

For more information and tips on coping with depression, see our publication [Depression after brain injury](#). You can also talk through your worries and get more information on your condition by calling the Headway helpline free on 0808 800 2244. Remember, you can also call the Samaritans 24 hours a day on 116 123.

### *Lack of insight*

After mild head injury, many people are unable to accurately judge their own actions and may behave inappropriately without realising it. This is fairly common in the first few days after injury when people may be uncharacteristically reckless and fail to recognise that they have any problems. Often people will insist on

Helpline: 0808 800 2244  
Email: [helpline@headway.org.uk](mailto:helpline@headway.org.uk)  
Web: [www.headway.org.uk](http://www.headway.org.uk)

returning to work or to driving before they are ready. Usually, after a week or two, people will suddenly realise what they have been doing and find it difficult to believe they could have acted that way.

On occasions, the lack of insight might persist for longer and may result in problems at home or at work, or in socially unacceptable behaviour. This can be very distressing for friends, family and workmates and is a very difficult problem to deal with. Generally the problem will resolve itself in time, but there are some things that family and friends can do to help:

- Provide information and education on this issue, for instance by sharing our publication [\*Insight and awareness after brain injury\*](#)
- Provide verbal feedback on behaviour – tell the person when they have behaved inappropriately, but do so in a discreet and non-confrontational way
- Meet others in a similar position – Headway groups and branches can offer family members support and a place to share their difficulties

---

## Section three: Other issues

### Alcohol

Don't expect alcohol to affect you in the usual way. A head injury can greatly reduce tolerance to alcohol so you should abstain as much as possible until you have fully recovered. Avoid alcohol completely in the first few days after injury. More information and guidance on this is available in our publication [\*Alcohol after brain injury\*](#).

### Compensation claims

If your injury was caused by an accident, assault or negligence, you may be entitled to compensation. It is important to consult a solicitor who specialises in head injury and Headway provides a list of approved personal injury solicitors at [www.headway.org.uk/supporting-you/in-your-area/head-injury-solicitors-directory](http://www.headway.org.uk/supporting-you/in-your-area/head-injury-solicitors-directory). You can find further information on claiming compensation in our publication [\*Claiming compensation after brain injury\*](#).

Helpline: 0808 800 2244  
Email: [helpline@headway.org.uk](mailto:helpline@headway.org.uk)  
Web: [www.headway.org.uk](http://www.headway.org.uk)

## Further accidents

It is important to note that whilst recovering, your reactions and co-ordination may be affected, so you should take care to avoid dangerous situations where possible in order to avoid further accidents. Sports players in particular who return to play too soon have a much higher rate of further injuries.

## Driving

Driving is a complex skill which we usually take for granted, but the effects of a mild head injury can impair reaction times and risk accidents. Try to avoid driving until your concentration and reactions are good enough, then start carefully and avoid long, stressful journeys and heavy traffic. In some cases it may be necessary to inform the licensing authority in your area of the UK about your injury, so speak to your GP about this. For further information on this subject see our publication [Driving after brain injury](#).

## Late diagnosis and misdiagnosis

If you are experiencing symptoms that you think are the result of a mild head injury that occurred some time ago, see your GP. It can be difficult to assess the effects of a mild head injury because:

- The injury may have happened a long time ago
- The effects are not always obvious
- It can be difficult to explain the symptoms to your GP
- The symptoms can be the same as many other conditions
- Even if a CT scan has been clear, there can still be persistent difficulties

For these reasons it would be useful to note down all your symptoms and take this publication with you to show your GP.

## Returning to work

It is important not to return to work before you are ready. You could try discussing the situation with your employer and explaining your symptoms to them. Showing

Helpline: 0808 800 2244  
Email: [helpline@headway.org.uk](mailto:helpline@headway.org.uk)  
Web: [www.headway.org.uk](http://www.headway.org.uk)



your employer this publication or our publication [Employer's guide to brain injury](#) may also help them to better understand the effects of your injury. You could then try to arrange a gradual return, with part-time hours and starting with light, routine duties.

It would be advisable to avoid making major decisions and becoming involved in stressful situations until you feel you are ready. This is especially the case in jobs with high levels of stress and pressure and where margins for error are small. Mistakes made because of the injury could damage your confidence and hinder your recovery.

For further information and guidance on this topic, see our publication [Returning to work after brain injury](#).

## **Social difficulties**

Any worries or problems you have will affect those around you. Often people who don't know about the effects of head injury won't realise that your difficulties are a result of your injury. It is important that, if you have had a mild head injury yourself, you show this booklet to friends and family so they will be able to understand how you are feeling and therefore be more able to help you.

You can also show friends and family Headway's series of relationships publications, such as [Partners' guide to brain injury](#) and [Friends' guide to brain injury](#), which include information about how the injury might have affected you and what they can do to help.

More general information on how relationships can be affected after brain injury is available in our publication [Relationships after brain injury](#).

## **Playing sports**

If you have had a mild head injury (often referred to as 'concussion' in the sporting world) and play contact sports, make sure you seek medical advice before continuing to play. Repeated mild head injuries can have cumulative effects and it is dangerous to expose yourself to the risk of a second injury whilst still recovering from the first one. All sports should have return to play protocols,

Helpline: 0808 800 2244  
Email: [helpline@headway.org.uk](mailto:helpline@headway.org.uk)  
Web: [www.headway.org.uk](http://www.headway.org.uk)

although some sports are more advanced than others in their work in this important area. If you are in any doubt about returning to play, speak to your coach or your GP, and don't feel rushed to return to contact sport until you are ready.

It is notoriously difficult to diagnose a concussion during a live sporting event, largely due to its often hidden symptoms and the fact that their presentation can often be delayed for several hours after the incident. It is therefore vital for players to take responsibility for any suspicion of concussion (for instance if they don't quite feel right, develop blurry vision or feel nauseous), and for fellow participants, coaches, officials and spectators to look out for signs as well. If a concussion is suspected, take the *if in doubt, sit it out!* approach.

For further information and resources on concussion management, see our publication [Concussion in sport](#).

**The booklet of this publication received a Commended Award at the British Medical Association Patient Information Awards 2010.**

**As a charity, we rely on donations from people like you to continue being able to provide free information to those affected by brain injury. To donate, or find out how else you can get involved with supporting our work, visit [www.headway.org.uk/get-involved](http://www.headway.org.uk/get-involved).**

**If you would like to leave feedback for this publication, please consider completing our short survey at [www.surveymonkey.co.uk/r/hwpublications](http://www.surveymonkey.co.uk/r/hwpublications) or contact us at [publications@headway.org.uk](mailto:publications@headway.org.uk).**

Last reviewed 2016. Next review 2025.

Helpline: 0808 800 2244  
Email: [helpline@headway.org.uk](mailto:helpline@headway.org.uk)  
Web: [www.headway.org.uk](http://www.headway.org.uk)